arXiv:1406.2520v1 [math.PR] 10 Jun 2014

On Decoupling of Functions of Normal Vectors II

Pavel G. Grigoriev and Stanislav A. Molchanov
Department of Mathematics and Statistics
University of North Carolina at Charlotte
Charlotte, NC 28223, USA
thepavel@mail.ru

Abstract

A decoupling type inequality for a sum of functions of Guassian vectors is established.
Key words: Decoupling, Gaussian vectors, Wick's polynomials, Hermite's polynomials.

1 Results

In [3] the following decoupling results were established.

Theorem 1.1. *For a normally distributed random vector* $\bar{Y} = (Y_i)_{i=1,\dots,d}$ *satisfying* $\mathsf{E}Y_i = 0$, $\mathsf{E}Y_i^2 = 1$, $i = 1, \dots, d$, *we have*

$$c^- \sum_{i=1}^d \|\varphi_i(Y_i)\|_2^2 \le \Big\| \sum_{i=1}^d \varphi_i(Y_i) \Big\|_2^2 \le c^+ \sum_{i=1}^d \|\varphi_i(Y_i)\|_2^2$$

for all measurable functions $\varphi_i : \mathbb{R} \to \mathbb{R}$ *satisfying* $\mathsf{E}\varphi_i(Y_i) = 0$ *with the constants* c^- *and* c^+ *being the smallest and the largest eigenvalues of the correlation matrix of* $\bar{Y}$. *Moreover the constants are the best possible.*

Theorem 1.2. *Let* $\bar{Y}_1 = (Y_{1,i})_{i=1,\dots,d_1}$ *and* $\bar{Y}_2 = (Y_{2,j})_{j=1,\dots,d_2}$ *be standard normal vectors with the correlations*

$$\mathsf{E}Y_{\alpha,i}Y_{\beta,j} = \begin{cases} 0, & \alpha = \beta, i \ne j \\ 1, & \alpha = \beta, i = j \\ \rho_{i,j}, & \alpha \ne \beta \end{cases}$$

Then

$$c_-\big(\|\varphi_1(\bar{Y}_1)\|_2^2 + \|\varphi_2(\bar{Y}_2)\|_2^2\big) \le \big\|\varphi_1(\bar{Y}_1) + \varphi_2(\bar{Y}_2)\big\|_2^2 \le c_+\big(\|\varphi_1(\bar{Y}_1)\|_2^2 + \|\varphi_2(\bar{Y}_2)\|_2^2\big)$$

holds for all measurable functions $\varphi_\alpha : \mathbb{R}^{d_\alpha} \to \mathbb{R}$ *such that* $\mathsf{E}\varphi_\alpha(\bar{Y}_\alpha) = 0$, $\alpha = 1, 2$, *with the constants* $c_\pm = 1 \pm s^*$, *where* s^* *is the maximum singular value of the matrix* $R = (\rho_{i,j})_{i=1,\dots,d_1, j=1,\dots,d_2}$. *These constants cannot be improved.*

These theorems refine the estimates used by Cherny et al. [1], [2]. Here we generalize these results further and prove

Theorem 1.3. *Let $\bar{Y}_\alpha = (Y_{\alpha,i})_{i=1,\dots,d_\alpha}$, $\alpha = 1,\dots,N$, be standard normal vectors with the correlations*

$$\mathsf{E} Y_{\alpha,i} Y_{\beta,j} = \begin{cases} 0, & \alpha = \beta, i \neq j \\ 1, & \alpha = \beta, i = j \\ \rho_{i,j}^{\alpha,\beta}, & \alpha \neq \beta \end{cases}$$

Then

$$C_- \sum_{\alpha=1}^N \|\varphi_\alpha(\bar{Y}_\alpha)\|_2^2 \le \Big\| \sum_{\alpha=1}^N \varphi_\alpha(\bar{Y}_\alpha) \Big\|_2^2 \le C_+ \sum_{\alpha=1}^N \|\varphi_\alpha(\bar{Y}_\alpha)\|_2^2 \tag{1}$$

for all measurable functions $\varphi_\alpha : \mathbb{R}^{d_\alpha} \to \mathbb{R}$, satisfying $\mathsf{E}\varphi_\alpha(\bar{Y}_\alpha) = 0$ for all $\alpha = 1,\dots,N$. The constants $C_\pm = 1 \pm \sigma_0$, where σ_0 denotes the largest eigenvalue of the matrix $S^* = (s^*_{\alpha,\beta})_{1\le\alpha,\beta\le N}$ with $s^*_{\alpha,\beta}$ being the maximum singular values of the matrices $R^{\alpha,\beta} = (\rho_{i,j}^{\alpha,\beta})_{i=1,\dots,d_\alpha, j=1,\dots,d_\beta}$ for $\alpha \neq \beta$ and $s^*_{\alpha,\alpha} := 0$.

Note that while Theorems 1.1 and 1.2 give the decoupling estimates with sharp constants, Theorem 1.3 provides just rough estimate with not the best constants (in particular C_- could be negative). However, in view of the applications described in Cherny et al. [1], [2] the upper bound in (1) is still interesting.

2 Proof of Theorem 1.3

The proof of Theorem 1.3 follows the framework used in [3]. We need to introduce notations used in [3] and borrowed from [4].

For $k = 0, 1, \dots$ we define Wick's polynomials $:x^k:$ by the extension

$$\exp\Big(ax - \frac{a}{2}\Big) = \sum_{k=0}^\infty a^k \frac{:x^k:}{k!}.$$

(Wick's polynomials are specially normalized Hermite's polynomials used in mathematical physics. We find these notations convenient for multidimensional case.)

Let $\bar{k} = (k_1,\dots,k_d)$ be a d-dimensional vector of non-negative integers. Set

$$\begin{aligned} |\bar{k}| &:= k_1 + \dots + k_d, \\ \bar{k}! &:= k_1! k_2! \dots k_d! \\ \bar{a}^{\bar{k}} &:= a_1^{k_1} a_2^{k_2} \dots a_d^{k_d}, \qquad \text{for } \bar{a} := (a_i)_{i=1,\dots,d} \in \mathbb{R}^d. \end{aligned}$$

For a vector variable $\bar{x} = (x_k)_{k=1}^d$ we define multidimensional Wick's polynomial by

$$:\bar{x}^{\bar{k}}: := \prod_{i=1}^d :x_i^{k_i}: .$$

It is well known (see e.g. [4]) that for a standard d-dimensional normal vector $\bar{Y}$ the system $\bigl\{\bar{k}!^{-1/2} : \bar{Y}^{\bar{k}} : \bigr\}_{\bar{k}\in\mathbb{Z}_0^d}$ is an orthonormal bases in the L_2 space generated by all square-integrable $f(\bar{Y})$. So for each $\alpha = 1, \dots, N$, we have

$$\varphi_\alpha(\bar{Y}_\alpha) = \sum_{\bar{k}\in\mathbb{Z}_0^{d_\alpha}} a_{\alpha,\bar{k}} \frac{: \bar{Y}_\alpha^{\bar{k}} :}{\bar{k}!^{1/2}} = \sum_{n=0}^{\infty} \sum_{\bar{k}\in\mathbb{Z}_0^{d_\alpha},\ |\bar{k}|=n} a_{\alpha,\bar{k}} \frac{: \bar{Y}_\alpha^{\bar{k}} :}{\bar{k}!^{1/2}}$$

and therefore

$$\sum_{\alpha=1}^{N} \varphi_\alpha(\bar{Y}_\alpha) = \sum_{\alpha=1}^{N} \sum_{n=0}^{\infty} \sum_{\bar{k}_\alpha\in\mathbb{Z}_0^{d_\alpha},\ |\bar{k}_\alpha|=n} a_{\alpha,\bar{k}_\alpha} \frac{: \bar{Y}_\alpha^{\bar{k}_\alpha} :}{\bar{k}_\alpha!^{1/2}} = \sum_{n=0}^{\infty} \sum_{\alpha=1}^{N} P_n(\bar{Y}_\alpha),$$

where $P_n(\bar{Y}_\alpha) := \sum\limits_{\bar{k}_\alpha\in\mathbb{Z}_0^{d_\alpha},\ |\bar{k}_\alpha|=n} a_{\alpha,\bar{k}_\alpha} \frac{:\bar{Y}_\alpha^{\bar{k}_\alpha}:}{\bar{k}_\alpha!^{1/2}}$. In particular,

$$\sum_{\alpha=1}^{N} \|\varphi_\alpha(\bar{Y}_\alpha)\|_2^2 = \sum_{n=0}^{\infty} \sum_{\alpha=1}^{N} \|P_n(\bar{Y}_\alpha)\|_2^2.$$

It is well-known that $P_{n_1}(\bar{Y}_\alpha)$ and $P_{n_2}(\bar{Y}_\beta)$ are orthogonal whenever $n_1 \neq n_2$ (see e.g. [3]). Consequently, to prove (1) for arbitrary φ_α it suffices to prove it for $\phi_\alpha(\bar{Y}_\alpha) = P_n(\bar{Y}_\alpha)$ for each $n = 0, 1, \dots$.

For a fixed n, we have

$$\Bigl\| \sum_{\alpha=1}^{N} P_n(\bar{Y}_\alpha) \Bigr\|_2^2 = \sum_{\alpha=1}^{N} \|P_n(\bar{Y}_\alpha)\|_2^2 + 2 \sum_{\alpha<\beta} \mathsf{E}\bigl[P_n(\bar{Y}_\alpha) P_n(\bar{Y}_\beta)\bigr]. \tag{2}$$

Fix a pair $\alpha < \beta$. Let $R^{\alpha,\beta} = U\Sigma V^T$ be the singular value decomposition of the matrix $R^{\alpha,\beta}$ (recall that here Σ is a $d_\alpha \times d_\beta$ diagonal matrix whose diagonal entries are the singular values of $R^{\alpha,\beta}$ and U, V are orthogonal matrices of corresponding sizes). Let $\bar{Z}_1 := U\bar{Y}_\alpha$ and $\bar{Z}_2 := V\bar{Y}_\beta$ (all the vectors are assumed being columns). This transformation can be written in the block matrix form

$$\begin{pmatrix} \bar{Z}_1 \\ \bar{Z}_2 \end{pmatrix} = \begin{pmatrix} U & 0 \\ 0 & V \end{pmatrix} \begin{pmatrix} \bar{Y}_\alpha \\ \bar{Y}_\beta \end{pmatrix}$$

with the obviously *orthogonal* $(d_\alpha + d_\beta) \times (d_\alpha + d_\beta)$ transformation matrix. It follows that

$$\mathsf{E}\bigl[P_n(\bar{Y}_\alpha) P_n(\bar{Y}_\beta)\bigr] = \mathsf{E}\bigl[P_n(\bar{Z}_1) P_n(\bar{Z}_2)\bigr] = \sum_{\substack{\bar{k}_\alpha\in\mathbb{Z}_0^{d_\alpha} \\ |\bar{k}_\alpha|=n}} \sum_{\substack{\bar{k}_\beta\in\mathbb{Z}_0^{d_\beta} \\ |\bar{k}_\beta|=n}} a_{\alpha,\bar{k}_\alpha} a_{\beta,\bar{k}_\beta} \mathsf{E} \frac{: \bar{Z}_1^{\bar{k}_\alpha} :}{\bar{k}_\alpha!^{1/2}} \frac{: \bar{Z}_2^{\bar{k}_\beta} :}{\bar{k}_\beta!^{1/2}} \tag{3}$$

Note that the covariance structure of $(\bar{Z}_1, \bar{Z}_2)$ is relatively simple, its covariance matrix is

$$\begin{pmatrix} I_{d_\alpha} & \Sigma \\ \Sigma^T & I_{d_\beta} \end{pmatrix}.$$

Without loss of generality we assume that $d_\alpha \geq d_\beta$, i.e. the vectors $\bar{Y}_\alpha$ are ordered according to their dimensions. Let us agree that the index vectors of different dimensions are equal ($\bar{k}_\alpha = \bar{k}_\beta$) if the shorter vector $\bar{k}_\beta$ coincides with the first d_β entries of k_α and the other entries of k_α are zeros. Also let us denote by $\bar{s}$ the vector of the diagonal entries of Σ (i.e. the singular values of $R^{\alpha,\beta}$).

Using Lemma 3.1 from [3] (for the case of diagonal R, see also formula (3.5) in [3]) we conclude

$$\mathsf{E} : \bar{Z}_1^{\bar{k}_\alpha} :: \bar{Z}_2^{\bar{k}_\beta} := \begin{cases} 0, & \bar{k}_\alpha \neq \bar{k}_\beta \\ \bar{k}_\beta! \bar{s}^{\bar{k}_\beta}, & \bar{k}_\alpha = \bar{k}_\beta \end{cases} \tag{4}$$

Using (4) we continue (3) as follows

$$\mathsf{E}\big[P_n(\bar{Y}_\alpha)P_n(\bar{Y}_\beta)\big] = \sum_{\substack{\bar{k}_\beta \in \mathbb{Z}_0^{d_\beta}, \bar{k}_\alpha \in \mathbb{Z}_0^{d_\alpha}: \\ |\bar{k}_\beta|=n,\, k_\alpha = \bar{k}_\beta}} a_{\alpha,\bar{k}_\alpha} a_{\beta,\bar{k}_\beta} \bar{s}^{\bar{k}_\beta}.$$

Taking into account that the singular values lie in $[0,1]$ (because they are certain correlations) we can estimate

$$\begin{aligned} \big|\mathsf{E}\big[P_n(\bar{Y}_\alpha)P_n(\bar{Y}_\beta)\big]\big| &\leq s^*_{\alpha,\beta} \sum_{\substack{\bar{k}_\beta \in \mathbb{Z}_0^{d_\beta}, \bar{k}_\alpha \in \mathbb{Z}_0^{d_\alpha}: \\ |\bar{k}_\beta|=n,\, k_\alpha = \bar{k}_\beta}} |a_{\alpha,\bar{k}_\alpha} a_{\beta,\bar{k}_\beta}| \\ &\leq s^*_{\alpha,\beta} \Big(\sum_{\substack{\bar{k}_\alpha \in \mathbb{Z}_0^{d_\alpha} \\ |\bar{k}_\alpha|=n}} a^2_{\alpha,\bar{k}_\alpha} \Big)^{\frac{1}{2}} \Big(\sum_{\substack{\bar{k}_\beta \in \mathbb{Z}_0^{d_\beta} \\ |\bar{k}_\beta|=n}} a^2_{\beta,\bar{k}_\beta} \Big)^{\frac{1}{2}} = s^*_{\alpha,\beta} \|P_n(\bar{Y}_\alpha)\|_2 \|P_n(\bar{Y}_\beta)\|_2 \end{aligned}$$

Recalling (2) we conclude

$$\Big| \Big\| \sum_{\alpha=1}^N P_n(\bar{Y}_\alpha) \Big\|_2^2 - \sum_{\alpha=1}^N \|P_n(\bar{Y}_\alpha)\|_2^2 \Big| \leq 2 \sum_{\alpha<\beta} s^*_{\alpha,\beta} \|P_n(\bar{Y}_\alpha)\|_2 \|P_n(\bar{Y}_\beta)\|_2.$$

By standard linear algebra argument we have that the right hand side is bounded by $\sigma_0 \sum_{\alpha=1}^N \|P_n(\bar{Y}_\alpha)\|_2^2$, where σ_0 is the largest eigenvalue of the matrix S^*. We conclude that for a fixed n the estimate (1) holds for $\phi_\alpha(\bar{Y}_\alpha) = P_n(\bar{Y}_\alpha)$, $\alpha = 1, \ldots, N$, with $C_\pm = 1 \pm \sigma_0$. As it was already pointed out this means that we have (1) for all ϕ_α with the same constants. $\square$

References

[1] *Cherny, A., Douady, R., Molchanov, S.* On measuring nonlinear risk with scarce observations. Finance Stoch. **14** (2010), no. 3, p. 375–395.

[2] *Cherny, A., Douady, R., Molchanov, S.* On Measuring Hedge Fund Risk. 2008. Preprint, available at: `http://mech.math.msu.su/~cherny`.

[3] *Grigoriev, P., Molchanov, S.* On Decoupling of Functions of Normal Vectors. Math. Notes **92** (2012), p. 401–409.

[4] *Malyshev, V.A., Minlos, R.A.* Gibbs Random Fields. Kluwer, Dordrecht (1991)